

LAMINATED CURRENTS

ABSTRACT. In this paper we prove the equivalence of two definitions of laminated currents.

JOHN ERIK FORNÆSS*, YINXIA WANG AND ERLEND FORNÆSS WOLD

1. INTRODUCTION

Let K be a relatively closed subset of the bidisc $\Delta^2(z, w) = \{(z, w); |z|, |w| < 1\}$. We suppose that K is a disjoint union of holomorphic graphs, $w = f_\alpha(z)$, where f_α is a holomorphic function on the unit disc with $f_\alpha(0) = \alpha$ and $|f_\alpha(z)| < 1$. We let $\mathcal{L}$ denote the lamination of K .

There are two notions of laminated currents that we will discuss, [1]. Let T be a positive closed $(1, 1)$ current supported on K . We assume that T is the restriction of a positive closed current defined on a neighborhood of $\overline{\Delta}^2$. We denote by $[V_\alpha]$ the current of integration along the graph of f_α . Let λ denote a continuous $(1, 0)$ form which at $(z, f_\alpha(z))$ equals a non zero multiple of $dw - f'_\alpha(z)dz$.

Definition 1. *We say that T is weakly directed by the lamination $\mathcal{L}$ if $\lambda \wedge T = 0$ for any such λ .*

Definition 2. *We say that T is directed by $\mathcal{L}$ if there is a positive measure μ so that $T = \int_\alpha [V_\alpha] d\mu(\alpha)$.*

Our main result is

Main Theorem. *The current T is directed if and only if it is weakly directed.*

We note that this is a result by Sullivan in the case of the lamination being smooth, i.e. the graphs vary smoothly with α , [7]. The part of Sullivan's proof that does not go through automatically in the non smooth case is a certain approximation step, and so in the present article we are concerned with approximation of partially smooth functions. In [2] the authors proved such an approximation theorem in the case of laminations in $\mathbb{R}^2$ and in $\mathbb{R}^3$.

In the last section we show that the main theorem breaks down for Riemann surface laminations in higher dimension.

2. PRELIMINARY ESTIMATES FOR SLOPES OF HOLOMORPHIC GRAPHS

The case we are studying is given by a holomorphic motion, see [8] for an exposition and further references. In this paper we need a basic estimate on slopes of the graphs. For the benefit of the reader we include the details of this well known fact.

*The first author is supported by an NSF grant. Keywords: Approximation, currents, test functions. 2000 AMS classification. Primary: 57R30, Secondary: 32U40

We start with a Lemma. Let $\Delta := \{z \in \mathbb{C}; |z| < 1\}$ denote the unit disc in $\mathbb{C}$. We denote by $\mathcal{O}(\Omega)$ the space of holomorphic functions on Ω . Let $\|\cdot\|_\infty$ denote the sup norm. Set

$$H^\infty = H^\infty(\Delta) = \{f \in \mathcal{O}(\Delta); \|f\|_\infty < \infty\}.$$

Also, if $0 < C < \infty$ we set

$$H_C^\infty = H_C^\infty(\Delta) = \{f \in \mathcal{O}(\Delta); \|f\|_\infty < C\}.$$

Lemma 1. *If $f \in H_1^\infty(\Delta)$ and $f(z) \neq 0 \forall z \in \Delta$, then $|f'(0)| \leq 2|f(0)| \log \frac{1}{|f(0)|}$.*

Proof. Pick a holomorphic function $f(z)$ on the unit disc such that $0 \neq |f(z)| < 1$ for all $z \in \Delta$. We can replace $f(z)$ by $e^{i\theta}f(z)$ for any real θ . This does not change $|f(0)|$ and $|f'(0)|$. Hence we can assume that $f(0) > 0$.

We set $h(z) := \log f(z)$. Then $h(z)$ is a holomorphic function on the unit disc and $\operatorname{Re}(h(z)) < 0$. We can also choose a branch of the logarithm so that $\log(f(0)) = -a < 0$. If $k(z) = \frac{h(z)}{a}$, then $k(z)$ is a holomorphic function on the unit disc and $k(0) = -1, \operatorname{Re}(k(z)) < 0$. We define $L(w) = \frac{w+1}{w-1}$. Then $L(-1) = 0$ and if $\operatorname{Re}(w) < 0$ then $|L(w)| < 1$. Then $\Gamma(z) := L(k(z))$ is a holomorphic function from the unit disc to the unit disc. Moreover $\Gamma(0) = L(k(0)) = L(-1) = 0$. Since $\Gamma(0) = 0$ and $|\Gamma(z)| < 1$ we can apply the Schwarz' Lemma. So we can conclude that $|\Gamma'(0)| \leq 1$. By the chain rule, $\Gamma'(0) = L'(k(0))k'(0) = L'(-1)k'(0)$. Since $L'(w) = \frac{-2}{(w-1)^2}$ we get $\Gamma'(0) = \frac{-2}{(-1-1)^2}k'(0)$ and therefore $k'(0) = -2\Gamma'(0)$. Hence we get $|k'(0)| \leq 2$. Since $k(z) = \frac{h(z)}{a}$, we next can conclude that $|k'(0)| = |h'(0)|/a$. Hence $|h'(0)| = a \cdot 2$ so $|h'(0)| \leq 2a$. Next recall that $h(z) = \log f(z)$ so $f(z) = e^{h(z)}$. Hence $f'(z) = e^{h(z)}h'(z)$. Therefore $f'(0) = e^{h(0)}h'(0) = f(0)h'(0)$. Hence $|f'(0)| \leq |f(0)||h'(0)|$. This implies that $|f'(0)| \leq 2a|f(0)|$. Now recall that $\log f(0) = -a$. But we have set this up so that $\log f(0) = \log |f(0)| + i \arg f(0)$ is real valued. So we have that $\log |f(0)| = -a$ i.e. $\log \frac{1}{|f(0)|} = a$. Therefore $|f'(0)| \leq 2a|f(0)| = 2|f(0)| \log \frac{1}{|f(0)|}$. This concludes the proof of the Lemma. $\square$

Corollary 1. *Suppose that we have two functions f and g holomorphic on the unit disk with $f - g \in H_1^\infty(\Delta)$. Suppose that $f(z) \neq g(z)$ for each $z \in \Delta$. We then have the estimate $|f'(z) - g'(z)| \leq 4|f(z) - g(z)| \log \frac{1}{|f(z) - g(z)|}$ for all $z \in \Delta, |z| < 1/2$.*

Proof. Pick $z, |z| < 1/2$. We define $G(w) = f(z + w/2) - g(z + w/2)$. Then $G(w)$ satisfies the conditions of Lemma 1. Hence $|G'(0)| \leq 2|G(0)| \log \frac{1}{|G(0)|}$. Therefore,

$$\frac{1}{2}|f'(z) - g'(z)| \leq 2|f(z) - g(z)| \log \frac{1}{|f(z) - g(z)|}$$

$\square$

3. APPROXIMATION FOR COMPLEX CURVES IN $\mathbb{C}^2$

We assume that for every $c = (a, b) = (a + ib) \in \mathbb{C}$ we have a holomorphic graph Γ_c given by $w = y_1 + iy_2 = f_c(z), z = x_1 + ix_2 \in \Delta$. We assume that all surfaces are disjoint and that there is a surface through every point in $\Delta \times \mathbb{C}$. We assume that $f_c(0) = c$.

Let $\pi : \Delta \times \mathbb{C} \rightarrow \mathbb{C}$ be defined by $\pi(z, f_c(z)) = c$. The lamination of $\Delta \times \mathbb{C}$ by the Γ_c 's defines a holomorphic motion and so by [5] the map $(z, c) \mapsto (z, f_c(z))$ is a continuous function in (z, c) . It follows that the function π is continuous.

Fix a positive constant R . By Corollary 1 there exists a positive real number $\delta_0 > 0$ such that if $z \in \frac{1}{2}\Delta$ and if $c, c' \in R\Delta$ with $|c - c'| < \delta_0$ then

$$(1) \left| \frac{\partial}{\partial z} f_{c'}(z) - \frac{\partial}{\partial z} f_c(z) \right| \leq 4 \cdot |f_{c'}(z) - f_c(z)| \log \frac{1}{|f_{c'}(z) - f_c(z)|}.$$

We define a class of partially smooth functions:

$$\begin{aligned} \mathcal{A} &:= \{ \phi \in \mathcal{C}(\Delta \times \mathbb{C}); \phi(z, f_c(z)) \in \mathcal{C}^1(\Gamma_c), \\ \Phi(x_1, x_2, w) &:= \frac{\partial}{\partial x_1} \phi(x_1, x_2, f_c(x_1, x_2)), w = f_c(x_1, x_2) \in \mathcal{C}(\Delta \times \mathbb{C}), \\ \Psi(x_1, x_2, w) &:= \frac{\partial}{\partial x_2} \phi(x_1, x_2, f_c(x_1, x_2)), w = f_c(x_1, x_2) \in \mathcal{C}(\Delta \times \mathbb{C}) \}. \end{aligned}$$

Theorem 1. *Let $\phi \in \mathcal{A}$, let R be a positive real number and let $\epsilon > 0$. Then there exists a function $\psi \in \mathcal{C}^1(\Delta \times R\Delta)$ such that for every point $(x_1, x_2, w) = (x_1, x_2, f_c(x_1, x_2)) \in \Delta \times R\Delta$:*

$$\begin{aligned} |\psi(x_1, x_2, w) - \phi(x_1, x_2, w)| &< \epsilon, \\ \left| \frac{\partial}{\partial x_1} [\psi(x_1, x_2, f_c(x_1, x_2))] - \frac{\partial}{\partial x_1} [\phi(x_1, x_2, f_c(x_1, x_2))] \right| &< \epsilon, \\ \left| \frac{\partial}{\partial x_2} [\psi(x_1, x_2, f_c(x_1, x_2))] - \frac{\partial}{\partial x_2} [\phi(x_1, x_2, f_c(x_1, x_2))] \right| &< \epsilon. \end{aligned}$$

We will prove the theorem using the following result:

Proposition 1. *Let $g \in \mathcal{A}$, $g(x_1, x_2, f_{a+ib}(x_1, x_2)) = a$, and let R be a positive real number. There exists a positive real number t_0 such that the following holds: For all $\epsilon > 0$ there exists a function $h \in \mathcal{C}^1(t_0\Delta \times R\Delta)$ such that for every point $(x_1, x_2, w) = (x_1, x_2, f_c(x_1, x_2)) \in t_0\Delta \times R\Delta$:*

$$\begin{aligned} |h(x_1, x_2, w) - g(x_1, x_2, w)| &< \epsilon, \\ \left| \frac{\partial}{\partial x_1} [h(x_1, x_2, f_c(x_1, x_2))] \right| &< \epsilon, \\ \left| \frac{\partial}{\partial x_2} [h(x_1, x_2, f_c(x_1, x_2))] \right| &< \epsilon. \end{aligned}$$

The same result holds if we replace a by b in the definition of g .

Proof of Theorem 1 from Proposition 1:

Lemma 2. *Let $p \in \Delta$ be a point, and let R, t_0 be positive real numbers such that $\Delta_{t_0}(p) \subset \subset \Delta$. Consider the lamination restricted to $\Delta_{t_0}(p) \times \mathbb{C}$. If the conclusion of Proposition 1 holds on $\Delta_{t_0}(p) \times R\Delta$ (with respect to projection onto $\{p\} \times \mathbb{C}$), then the conclusion of Theorem 1 holds on $\Delta_{t_0}(p) \times R\Delta$.*

Proof. Let $\pi = (\pi_1, \pi_2)$ denote the projection onto $\{p\} \times \mathbb{C}$. For each $j, k \in \mathbb{Z}$ and $\delta > 0$ we let $c^\delta(j, k)$ denote the point $(p, j\delta + k\delta i)$. Let Λ_j^δ denote the $\mathcal{C}^1$ -smooth function defined by $\Lambda_j^\delta(t) = \cos^2[\frac{\pi}{2\delta}(t - j\delta)]$ when $(j-1)\delta \leq t \leq (j+1)\delta$ and 0 otherwise. For each $c^\delta(j, k)$ we first define a function

$$\psi_{jk}^\delta(z) := \phi(z, f_{c^\delta(j,k)}(z)),$$

and then we define a preliminary approximation

$$\psi^\delta(z, w) = \sum_{j,k} \psi_{jk}^\delta(z) \Lambda_j(\pi_1(z, w)) \Lambda_k(\pi_2(z, w)).$$

Let $(z_0, w_0) \in \triangle_{t_0}(p) \times R\triangle$. Then $\pi(z_0, w_0)$ is contained in a square with corners $c^\delta(j, k), c^\delta(j+1, k), c^\delta(j, k+1)$ and $c^\delta(j+1, k+1)$, and we have that

$$\psi^\delta(z_0, w_0) = \sum_{m=j, j+1, n=k, k+1} \psi_{mn}^\delta(z_0) \Lambda_m(\pi_1(z_0, w_0)) \Lambda_n(\pi_2(z_0, w_0)).$$

We have that

$$\begin{aligned} & |\psi^\delta(z_0, w_0) - \phi(z_0, w_0)| \\ &= \left| \sum_{m=j, j+1, n=k, k+1} [\psi_{mn}^\delta(z_0) - \phi(z_0, w_0)] \cdot \Lambda_m^\delta(\pi_1(z_0, w_0)) \cdot \Lambda_n^\delta(\pi_2(z_0, w_0)) \right| \\ &\leq \max_{m=j, j+1, n=k, k+1} \{ |\psi_{mn}^\delta(z_0) - \phi(z_0, w_0)| \} \end{aligned}$$

Since the map from $\overline{\triangle}_{t_0}(p) \times \mathbb{C}$ defined by $(z, \alpha) \mapsto (z, f_\alpha(z))$ is a homeomorphism it follows that $\psi^\delta \rightarrow \phi$ uniformly as $\delta \rightarrow 0$.

Next we approximate derivatives along leaves. Let α be such that $(z_0, w_0) = (z_0, f_\alpha(z_0))$. Since the functions $\Lambda_j^\delta \circ \pi_i$ are constant along leaves we get that

$$\begin{aligned} & \left| \frac{\partial}{\partial x_i} [\psi^\delta(z_0, f_\alpha(z_0)) - \phi(z_0, f_\alpha(z_0))] \right| \\ &= \left| \sum_{m=j, j+1, n=k, k+1} \left[\frac{\partial}{\partial x_i} [\psi_{mn}^\delta(z_0) - \phi(z_0, f_\alpha(z_0))] \right] \right. \\ &\quad \times \left. \Lambda_m^\delta(\pi_1(z_0, f_\alpha(z_0))) \cdot \Lambda_n^\delta(\pi_2(z_0, f_\alpha(z_0))) \right| \\ &\leq \max_{m=j, j+1, n=k, k+1} \left\{ \left| \frac{\partial}{\partial x_i} [\psi_{mn}^\delta(z_0) - \phi(z_0, f_\alpha(z_0))] \right| \right\} \end{aligned}$$

It follows that $\psi^\delta \rightarrow \phi$ also in $\mathcal{C}^1$ -norm on leaves.

Now the conclusion of Lemma 2 follows because the functions π_j can be approximated uniformly and in $\mathcal{C}^1$ -norm on leaves. $\square$

For each point $p \in \triangle$ there exists by Proposition 1 a positive real number t_p such that constant approximation is possible on $\triangle_{t_p}(p) \times R\triangle$. Hence by Lemma 2 approximation of functions in $\mathcal{A}$ is possible.

We may then choose a locally finite cover $\{U_\alpha\}_{\alpha \in \mathbb{N}}$ of $\triangle$ by disks such that approximation by functions in $\mathcal{A}$ is possible on each $U_\alpha \times R\triangle$. Let $\{\varphi_\alpha\}$ be a partition of unity subordinate to $\{U_\alpha\}$. For each α let $C_\alpha = \|\nabla \varphi_\alpha\|$.

For a given ϵ_α let g_{ϵ_α} be an ϵ_α -approximating function of ϕ on $U_\alpha \times R\triangle$. We will show that there is a sequence $\{\epsilon_\alpha\}$ such that the function

$$\psi = \sum_{\alpha} \varphi_\alpha \cdot g_{\epsilon_\alpha}$$

satisfies the claims of the Theorem.

Let $(z_0, f_c(z_0)) \in U_\alpha$, and let $\alpha_1, \dots, \alpha_m$ be the finite number of α_i 's such that the support of ϕ_α intersects U_α . Then

$$\psi(z, f_c(z)) = \sum_{i=1}^m \varphi_{\alpha_i}(z) \cdot g_{\epsilon_{\alpha_i}}(z, f_c(z))$$

for all z near z_0 . Then

$$\begin{aligned} |\psi(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))| &= \left| \left[\sum_{i=1}^m \varphi_{\alpha_i}(z_0) \cdot g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) \right] - \phi(z_0, f_c(z_0)) \right| \\ &\leq \sum_{i=1}^m \varphi_{\alpha_i}(z_0) \cdot |g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))| \\ &\leq \max\{\epsilon_{\alpha_i}\} \end{aligned}$$

Further

$$\begin{aligned} & \left| \frac{\partial}{\partial x_1} [\psi(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))] \right| \\ &= \left| \frac{\partial}{\partial x_1} \left[\left[\sum_{i=1}^m \varphi_{\alpha_i}(z) \cdot g_{\epsilon_{\alpha_i}}(z, f_c(z_0)) \right] - \phi(z_0, f_c(z_0)) \right] \right| \\ &= \left| \sum_{i=1}^m \frac{\partial}{\partial x_1} [\varphi_{\alpha_i}(z_0) \cdot (g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0)))] \right| \\ &= \left| \sum_{i=1}^m \frac{\partial}{\partial x_1} [\varphi_{\alpha_i}(z_0)] \cdot (g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))) \right| \\ &+ \sum_{i=1}^m \varphi_{\alpha_i}(z_0) \cdot \left| \frac{\partial}{\partial x_1} [g_{\epsilon_{\alpha_i}}(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))] \right| \\ &\leq m \cdot \max\{C_{\alpha_i}\} \cdot \max\{\epsilon_{\alpha_i}\} + \max\{\epsilon_{\alpha_i}\} \end{aligned}$$

Similarly we get that

$$\left| \frac{\partial}{\partial x_2} [\psi(z_0, f_c(z_0)) - \phi(z_0, f_c(z_0))] \right| \leq m \cdot \max\{C_{\alpha_i}\} \cdot \max\{\epsilon_{\alpha_i}\} + \max\{\epsilon_{\alpha_i}\}$$

It is clear that we may choose ϵ_{α_i} for $i = 1, \dots, m$ to get the desired estimate for all points $z_0 \in U_\alpha$ for this particular α . Running through all α 's we have that any particular α_i will only come under consideration a finite number of times. Hence we may choose the sequence $\{\epsilon_\alpha\}$. $\square$

We proceed to prove the Proposition.

Fix δ_0 to get the estimate (1) (in the beginning of Section 3) for all $|c - c'| < \delta_0$ with $|c|, |c'| \leq 2R$. For any δ with $0 < \delta < \delta_0$ we let $c^\delta(j, k) = (j + k \cdot i) \cdot \delta$ for $j, k \in \mathbb{Z}$. Let $\chi : [0, 1] \rightarrow \mathbb{R}$ be a smooth function such that $\chi(t) = 0$ for $0 \leq t \leq \frac{1}{4}$ and $\chi(t) = 1$ for $\frac{3}{4} \leq t \leq 1$. Let C be a constant such that $|\chi'(t)| \leq C$ for all $t \in [0, 1]$.

We first define a function h_δ on the surfaces $\Gamma_{c^\delta(j,k)}$ simply by $h_\delta|_{\Gamma_{c^\delta(j,k)}} \equiv j\delta$. We want to interpolate this function between the surfaces.

For a fixed z consider the sets of points

$$\Gamma_{c^\delta(j,k)}(z) := \{f_{c^\delta(j,k)}(z), f_{c^\delta(j+1,k)}(z), f_{c^\delta(j,k+1)}(z), f_{c^\delta(j+1,k+1)}(z)\}.$$

We first show that these sets move nicely with z for small enough $|z|$ and independent of δ . In particular we want to know that we may define quadrilateral regions $R_{\delta,j,k}(z)$ with corners $\Gamma_{c^\delta(j,k)}(z)$ and that these sets have disjoint interior.

We make the change of coordinates in the w variable, by setting

$$\tilde{w}(z, w) = \tilde{w}_{jk}(zw) = \frac{w - f_{c^\delta(j,k)}(z)}{f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)}.$$

We get that

$$\begin{aligned} \tilde{w}(z, f_{c^\delta(j,k)}(z)) &\equiv 0 \\ \tilde{w}(z, f_{c^\delta(j+1,k)}(z)) &\equiv 1 \end{aligned}$$

Lemma 3. *Fix N . Then there exists a real number $t_0 > 0$ independent of δ so that if $|l|, |m| < N$ then $|\tilde{w}_{jk}(z, f_{c^\delta(j+l,k+m)}(z)) - \tilde{w}_{jk}(z, f_{c^\delta(j+l,k+m)}(0))| < 1/10$ for all $|z| < t_0$ and any j, k .*

Proof. Let $\pi : \Delta \rightarrow \mathbb{C} \setminus \{0, 1\}$ be the universal cover, and fix a pair (j, k) . We have that $p_{lm}^\delta := \tilde{w}_{jk}(0, f_{c^\delta(j+l,k+m)}(0)) = l + m \cdot i$ for all δ , so we may choose a point $P_{lm} \in \Delta$ such that $\pi(P_{lm}) = p_{lm}^\delta$ for all δ . For each δ, j, k, l, m we have that $\tilde{w}_{jk}(z, f_{c^\delta(j+l,k+m)}(z))$ is a map $k^\delta : \Delta \rightarrow \mathbb{C} \setminus \{0, 1\}$, and so they lift to maps $g^\delta : \Delta \rightarrow \Delta$ with $g^\delta(0) = P_{lm}$, i.e. $k^\delta = \pi \circ g^\delta$. By the Schwartz' Lemma we have that $|g^\delta(z) - P_{lm}| \leq L_{lm}|z|$ for all δ . So if $|z|$ is small enough we have that $g^\delta(z) \subset \pi^{-1}(\Delta_{\frac{1}{10}}(p_{lm}))$. Since there are only a finite number of pairs (l, m) bounded by N the result follows. $\square$

From now on we assume that $|z| \leq t_0$.

Lemma 4. *The quadrilaterals have disjoint interiors.*

Proof. Pick (j, k) . We use the linear change of coordinates in the w direction for fixed z :

$$\tilde{w}_{jk}(z, w) = \frac{w - f_{c^\delta(j,k)}(z)}{f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)}.$$

This sends $f_{c^\delta(j+l,k+m)}(z)$ close to $(j+l, k+m)$ on a small disc in the z direction for uniformly bounded (l, m) . Hence it is clear that the quadrilaterals are disjoint. $\square$

Next we define preliminary functions h_{jk}^δ on the respective quadrilaterals. First we define a function $t_z(y_1, y_2)$ to be constant equal to 0 on the line between $f_{c^\delta(j,k)}(z)$ and $f_{c^\delta(j,k+1)}(z)$, and constant equal to 1 on the line between $f_{c^\delta(j+1,k)}(z)$ and $f_{c^\delta(j+1,k+1)}(z)$. We extend t_z continuously to be affine on the two other edges, and then we extend t_z to be constant equal to v on the line between $f_{c^\delta(j,k)}(z) + v \cdot (f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z))$ and $f_{c^\delta(j,k+1)}(z) + v \cdot (f_{c^\delta(j+1,k+1)}(z) - f_{c^\delta(j,k+1)}(z))$. Finally we define h_{jk}^δ by

$$h_{jk}^\delta(z, y_1, y_2) = j\delta + \delta \cdot (\chi \circ t_z)(y_1, y_2).$$

The h_{jk}^δ 's patch up smoothly in the "vertical" directions where the functions are constant. To be able to patch them together in the "horizontal" directions we first extend each h_{jk}^δ across the "horizontal" edges.

To do this we use the coordinates defined by $\tilde{w}$. Consider the normalization

$$\tilde{w}_{jk}(z, w) = \frac{w - f_{c^\delta(j,k)}(z)}{f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)}$$

Let $\tilde{h}_{jk}^\delta$ be defined by $\tilde{h}_{jk}^\delta \circ \tilde{\omega} = h_{jk}^\delta$. We want to glue together the two functions on the quadrilaterals sharing (in the new coordinates) the line segment γ between $(0, 0)$ and $(1, 0)$, i.e. the function $\tilde{h}_{jk}^\delta$ defined above γ and the function $\tilde{h}_{j(k-1)}^\delta$ below γ .

We start by extending the function $\tilde{h}_{jk}^\delta$. Note first that by Lemma 3 the quadrilaterals $R_{\delta,j,k}$ and $R_{\delta,j,k-1}$ in the new coordinates - henceforth denoted $\tilde{R}_{\delta,j,k}$ and $\tilde{R}_{\delta,j,k-1}$ - have corners within $\frac{1}{10}$ -distance from the points (l, m) for $l, m \in \{0, 1, -1\}$. Note also that if we define a function $\tilde{t}_z(\tilde{y}_1, \tilde{y}_2)$ ($\tilde{w} = \tilde{y}_1 + i\tilde{y}_2$) along lines in the quadrilateral $\tilde{R}_{\delta,j,k}(z)$ in the new coordinates as we did when we defined $t_z(y_1, y_2)$ above, then $h_{jk}^\delta = (j\delta + \delta(\chi \circ \tilde{t})) \circ \tilde{w}$. Because of the placing of the corners we see that there exists a constant K independent of δ, j, k such that $\|\nabla_{\tilde{w}}(j\delta + \delta(\chi \circ \tilde{t}))\| \leq K\delta$.

Continue the lines in $\tilde{R}_{\delta,j,k}$ that pass through the interval $[\frac{1}{8}, 1 - \frac{1}{8}]$ and extend $\tilde{h}_{jk}^\delta$ to be constant on these lines. By the placing of the corners there is a constant μ - independent of δ and j, k - such that these lines can be extended to the line between $(0, -\mu)$ and $(1, -\mu)$. Let $\tilde{P}_{\delta,j,k}$ denote the extended set $\tilde{R}_{\delta,j,k} \cup (\tilde{R}_{\delta,j,k-1} \cap \{y_2 \geq -\mu\})$, and we see that $\tilde{h}_{jk}^\delta$ extends to be constant on the part of $\tilde{P}_{\delta,j,k}$ where it is not already defined. Extend $\tilde{h}_{j(k-1)}^\delta$ similarly in the other direction.

To glue the functions together we choose a smooth function $\varphi(z, \tilde{y}_1, \tilde{y}_2) = \varphi(\tilde{y}_2)$ such that $\varphi(\tilde{y}_2) = 1$ if $y_2 \geq \mu$ and such that $\varphi(\tilde{y}_2) = 0$ if $y_2 \leq -\mu$. We define

$$h_\delta(z, w) := (\varphi \circ \tilde{w}_{jk})(z, w) \cdot h_{jk}^\delta(z, w) + (1 - \varphi \circ \tilde{w}_{jk})(z, w) \cdot h_{j(k-1)}^\delta(z, w)$$

Fix a constant M such that $\|\frac{\partial \varphi}{\partial \tilde{y}_2}\| = M$.

Lemma 5. *There are constants N_1 and N_2 such that for each j, k, δ we have that $h_{jk}^\delta(z, w) = j\delta$ if $|w - f_{c^\delta(j,k)}(z)| \leq N_1 |f_{c^\delta(j+1,k)}(z) - f_{c^\delta(j,k)}(z)|$. Moreover there is a smooth function $\tilde{g}_{jk}^\delta(z, \tilde{y}_1, \tilde{y}_2)$ such that $h_{jk}^\delta = \tilde{g}_{jk}^\delta \circ \tilde{w}$ and such that $\|\nabla_{\tilde{w}} \tilde{g}_{jk}^\delta\| \leq N_2 \delta$.*

Proof. The existence of the constant N_1 can be seen by our description of the function in local coordinates where we used Lemma 3. To see the rest let us give the function $\tilde{g}_{jk}^\delta$ explicitly.

Fix z . Let (a_1, a_2) denote the corner of $\tilde{R}_{j,k}^\delta$ which is close to $(0, 1)$, and define a map $A_z(\tilde{y}_1, \tilde{y}_2) := (\tilde{y}_1 - \tilde{y}_2 \frac{a_1}{a_2}, \tilde{y}_2 \frac{1}{a_2})$. Then A_z changes smoothly with z and we have that $\|A_z\| < 2$ for all the possibilities of (a_1, a_2) we are considering.

Next we define a function $\hat{t}$ on the quadrilateral $A_z(\tilde{R}_{j,k}^\delta)$ along lines as above. Let (b_1, b_2) denote the corner close to $(1, 1)$ and fix $\hat{y} = (\hat{y}_1, \hat{y}_2)$. We have that the two vertical sides of $A_z(\tilde{R}_{j,k}^\delta)$ meet at the point $(0, -L)$ where L is given by $L = \frac{b_2}{b_1 - 1}$. Calculating the slope of the line from the point $\hat{y}$ to the point $(\hat{t}(\hat{y}), 0)$

we get that $\frac{\widehat{y}_1}{L+\widehat{y}_2} = \frac{\widehat{t}(y)}{L}$ which gives us

$$\widehat{t}(\widehat{y}) = \frac{\widehat{y}_1 \cdot L}{L + \widehat{y}_2} = \frac{\widehat{y}_1 - b_2}{b_2 + \widehat{y}_2(b_1 - 1)}.$$

We have that $\widehat{t}$ varies smoothly with (b_1, b_2) and we see that $\widehat{t}$ has bounded derivatives for the cases of (b_1, b_2) we are considering. Define $\tilde{g}_{jk}^\delta$ by

$$\tilde{g}_{jk}^\delta = j\delta + \delta(\chi \circ \widehat{t} \circ A_z),$$

and the function h_{jk}^δ is given by $h_{jk}^\delta = \tilde{g}_{jk}^\delta \circ \tilde{w}$. $\square$

Lemma 6. $h_\delta \rightarrow g$ in sup norm on $\triangle_{t_0} \times R\triangle$.

Proof. It is clear that $h_\delta(0, \cdot) \rightarrow g(0, \cdot)$ uniformly. The claim then follows from Lemma 8 below. $\square$

Lemma 7. If t_0 and δ is small enough we have that $|f_{c^\delta(j,k)}(z) - f_{c^\delta(j+1,k)}(z)| \geq \delta^2$ for all z with $|z| \leq t_0$ and all j, k such that $|c^\delta(j, k)| \leq 2R$.

Proof. If δ is small enough we have the estimate

$$|f'_{c^\delta(j,k)}(z) - f'_{c^\delta(j+1,k)}(z)| \leq 4|f_{c^\delta(j,k)}(z) - f_{c^\delta(j+1,k)}(z)| \log \frac{1}{|f_{c^\delta(j,k)}(z) - f_{c^\delta(j+1,k)}(z)|}$$

for all $|z| \leq t_0$ and all $|c^\delta(j, k)| \leq 2R$ (Corollary 1). Let ϕ denote the function

$$\phi(z) = |f_{c^\delta(j,k)}(z) - f_{c^\delta(j+1,k)}(z)|,$$

and let $\phi(t)$ denote the restriction to radial real lines starting at the origin. We have that

$$\frac{\partial}{\partial t}[-\log \phi(t)] = -\frac{\phi'(t)}{\phi(t)},$$

and so

$$|\frac{\partial}{\partial t}[-\log \phi(t)]| \leq 4|-\log \phi(t)|.$$

It follows that

$$-\log \phi(t) \leq -\log \phi(0) \cdot e^{4t} \Rightarrow \phi(t) \geq \phi(0)e^{4t},$$

and since $\phi(0) = \delta$ the result follows by choosing t_0 smaller than $\frac{1}{4} \log 2$. $\square$

Lemma 8. Let $c = a + ib, j\delta \leq a \leq (j+1)\delta, k\delta \leq b \leq (k+1)\delta$. The function $h_\delta(z, f_c(z))$ is small in C^1 norm along the graph Γ_c .

Proof. We need to estimate the derivatives of the function $h_\delta(z, f_c(z))$ at an arbitrary point $(z_0, f_c(z_0))$. We estimate $\frac{\partial}{\partial x} = \frac{\partial}{\partial x_1}$ - the case of $\frac{\partial}{\partial x_2}$ is similar. Since we are working on lines we use the notation (x, y_1, y_2) for coordinates.

We observe first that if $(z_0, f_c(z_0))$ is outside $R_{\delta,j,k}$ then it must still be very close. If the point is close to the vertical edges, then the function h_δ is locally constant, so we are done. We can assume that also $(z_0, f_c(z_0)) \in P_{\delta,j,k} \setminus P_{\delta,j,k+1}$. We divide the proof into two cases: Assume first that $(z_0, f_c(z_0))$ is not in $R_{\delta,j,k-1}$. Then the function h_δ is simply equal to the function h_{jk}^δ .

We have that

$$\begin{aligned}
\frac{\partial}{\partial x}(h_{jk}^\delta(x, f(x))) &= \left(\frac{\partial h_{jk}^\delta}{\partial x}, \frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2}\right)(x, f(x)) \cdot \left(1, \frac{\partial f_1}{\partial x}, \frac{\partial f_2}{\partial x}\right)(x) \\
&= \frac{\partial h_{jk}^\delta}{\partial x}(x, f(x)) + \left(\frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2}\right)(x, f(x)) \cdot \left(\frac{\partial f_1}{\partial x}, \frac{\partial f_2}{\partial x}\right)(x)
\end{aligned}$$

For fixed s, v we may define a curve $(x, g(x))$:

$$\begin{aligned}
g(x) &= (1-s)[(1-v)f_{c^\delta(j,k)}(x) + vf_{c^\delta(j+1,k)}(x)] \\
&\quad + s[(1-v)f_{c^\delta(j,k+1)}(x) + vf_{c^\delta(j+1,k+1)}(x)].
\end{aligned}$$

Then $h_{jk}^\delta(x, g(x)) \equiv j\delta + \chi(v)\delta$. Choose s and v so that $(x_0, g(x_0)) = (x_0, f_c(x_0))$. We get that

$$0 = \frac{\partial}{\partial x}(h_{jk}^\delta(x, g(x))) = \frac{\partial h_{jk}^\delta}{\partial x}(x, g(x)) + \left(\frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2}\right)(x, g(x)) \cdot \left(\frac{\partial g_1}{\partial x}, \frac{\partial g_2}{\partial x}\right)(x),$$

and so

$$\frac{\partial}{\partial x}(h_{jk}^\delta(x_0, f(x_0))) = \left(\frac{\partial h_{jk}^\delta}{\partial y_1}, \frac{\partial h_{jk}^\delta}{\partial y_2}\right)(x_0, g(x_0)) \cdot \left(\frac{\partial f_1}{\partial x} - \frac{\partial g_1}{\partial x}, \frac{\partial f_2}{\partial x} - \frac{\partial g_2}{\partial x}\right)(x_0).$$

Using the Lemma 3 we see that $\|f_c(x_0) - f_{c^\delta(j+l,k+m)}(x_0)\| \leq 2\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|$ for $l, m \in \{0, 1\}$, and so

$$\begin{aligned}
&\left\| \frac{\partial}{\partial x}(f_c - f_{c^\delta(j+l,k+m)})(x_0) \right\| \\
&\leq 4\|(f_c - f_{c^\delta(j+l,k+m)})(x_0)\| \log \frac{1}{\|(f_c - f_{c^\delta(j+l,k+m)})(x_0)\|} \\
&\leq 8\|(f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)})(x_0)\| \log \frac{1}{2\|(f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)})(x_0)\|}.
\end{aligned}$$

It follows that

$$\begin{aligned}
&\left\| \frac{\partial}{\partial x}(h_\delta(x, f(x))) \right\| \\
&\leq 8 \cdot \left\| \left(\frac{\partial h_\delta}{\partial y_1}, \frac{\partial h_\delta}{\partial y_2}\right) \right\| \cdot \|(f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)})(x_0)\| \\
&\quad \times \log \frac{1}{2\|(f_{c^\delta(j+1,k)} - f_{c^\delta(j,k)})(x_0)\|}.
\end{aligned}$$

We proceed to estimate $\left\| \left(\frac{\partial h_\delta}{\partial y_1}, \frac{\partial h_\delta}{\partial y_2}\right) \right\|$. We change coordinates according to Lemma 5 and write h_δ as a composition $\tilde{g}_\delta \circ \tilde{w}(y)$. We get $\|D_w \tilde{w}\| = \frac{1}{\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|}$, and we have that $\|\nabla_{\tilde{w}} \tilde{g}_\delta\| \leq N_2 \delta$. This shows that

$$\left\| \left(\frac{\partial h_\delta}{\partial y_1}, \frac{\partial h_\delta}{\partial y_2}\right) \right\| \leq N_2 \delta \frac{1}{\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|}.$$

This gives

$$\left\| \frac{\partial}{\partial x}(h_\delta(x, f(x))) \right\| \leq 8N_2\delta \log \frac{1}{\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\|}.$$

We have by Lemma 7 that $\|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)\| \geq \delta^2$ and so

$$\left\| \frac{\partial}{\partial x}(h_\delta(x_0, f(x_0))) \right\| \leq 8N_2\delta \log \frac{1}{2\delta^2} \rightarrow 0 \text{ as } \delta \rightarrow 0.$$

The other case we have to consider is when $(z_0, f_c(z_0))$ is contained in an overlap where we glued our functions together. In that case we may assume that $(z_0, f_c(z_0))$ is also contained in $P_{j(k-1)}^\delta$.

Let $\vec{v}$ denote the vector $\vec{v} = \frac{\partial}{\partial x}(x_0, f_c(x_0))$. We have that

$$\begin{aligned} \nabla h_\delta(x_0, f_c(x_0)) \cdot \vec{v} &= \nabla[\varphi \circ \tilde{w} \cdot h_{jk}^\delta](x_0, f_c(x_0)) \cdot \vec{v} \\ &+ \nabla[(1 - \varphi) \circ \tilde{w} \cdot h_{j(k-1)}^\delta](x_0, f_c(x_0)) \cdot \vec{v} \\ &= h_{jk}^\delta(x_0, f_c(x_0)) \cdot \nabla[\varphi \circ \tilde{w}](x, f_c(x_0)) \cdot \vec{v} \\ &+ (\varphi \circ \tilde{w})(x_0, f_c(x_0)) \cdot \nabla[h_{jk}^\delta](x_0, f_c(x_0)) \cdot \vec{v} \\ &+ h_{j(k-1)}^\delta(x_0, f_c(x_0)) \cdot \nabla[(1 - \varphi) \circ \tilde{w}](x, f_c(x_0)) \cdot \vec{v} \\ &+ ((1 - \varphi) \circ \tilde{w})(x_0, f_c(x_0)) \cdot \nabla[h_{j(k-1)}^\delta](x_0, f_c(x_0)) \cdot \vec{v} \end{aligned}$$

By the above calculations we need not worry about the second and fourth term in this sum so we have to check that

$$(h_{jk}^\delta(x_0, f_c(x_0)) - h_{j(k-1)}^\delta(x_0, f_c(x_0))) \cdot \nabla[\varphi \circ \tilde{w}](x_0, f_c(x_0)) \cdot \vec{v} \rightarrow 0$$

as $\delta \rightarrow 0$.

First of all we have that $|h_{jk}^\delta(x_0, f_c(x_0)) - h_{j(k-1)}^\delta(x_0, f_c(x_0))| \leq 2\delta$. Further $|\nabla[\varphi \circ \tilde{w}](x_0, f_c(x_0)) \cdot \vec{v}| \leq M \cdot \|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\|$.

Now

$$D[\tilde{w}](x_0, f_c(x_0))(\vec{v}) = \frac{\partial}{\partial x} \left[(x, \frac{f_c(x) - f_{c^\delta(j,k)}(x)}{f_{c^\delta(j+1,k)}(x) - f_{c^\delta(j,k)}(x)}) \right](x_0).$$

Ignoring the constant term (it gets killed by δ) we get that

$$\begin{aligned}
& \|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\| \\
& \leq \frac{|f'_c(x_0) - f'_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \\
& + \frac{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)| \cdot |f'_{c^\delta(j+1,k)}(x_0) - f'_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|^2} \\
& \leq \frac{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \log \frac{1}{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|} \\
& + \frac{|(f_c(x_0) - f_{c^\delta(j,k)}(x_0))| \cdot |(f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0))|}{|(f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0))|^2} \\
& \times \log \frac{1}{|(f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0))|}
\end{aligned}$$

By Lemma 3 we have that $\frac{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \leq 2$ and so

$$\begin{aligned}
\|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\| & \leq 2 \cdot \log \frac{1}{|f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)|} \\
& + 2 \log \frac{1}{|f_c(x_0) - f_{c^\delta(j,k)}(x_0)|}
\end{aligned}$$

By Lemma 5 we have that our function is constant unless $|f_c(x_0) - f_{c^\delta(j,k)}(x_0)| \geq N_1 |f_{c^\delta(j+1,k)}(x_0) - f_{c^\delta(j,k)}(x_0)| \geq N_1 \delta^2$ (by Lemma 7), and so we may assume that

$$\|D[\tilde{w}](x_0, f_c(x_0))(\vec{v})\| \leq 2 \log \frac{1}{\delta^2} + 2 \log \frac{1}{N_1 \delta^2}.$$

All in all:

$$\begin{aligned}
& |(h_{jk}^\delta(x_0, f_c(x_0)) - h_{j(k-1)}^\delta(x_0, f_c(x_0))) \cdot \nabla[\varphi \circ \tilde{w}](x_0, f_c(x_0)) \cdot \vec{v}| \\
& \leq 4M\delta(\log \frac{1}{\delta^2} + \log \frac{1}{N_1 \delta^2}) \rightarrow 0 \text{ as } \delta \rightarrow 0.
\end{aligned}$$

□

4. PROOF OF THE MAIN THEOREM

We are ready to prove the main theorem. By the theorem of Slodkowski, [6], [8], we can assume that $\mathcal{L}$ is a lamination of $\Delta \times \mathbb{C}$ as in the previous section.

Proof of the Main Theorem: Suppose that T is a positive closed $(1, 1)$ current on $\Delta^2(0, 1)$, supported on the laminated set K described in the introduction. We assume that T is directed by the lamination $\mathcal{L}$ of K . Hence there is a positive measure μ so that $T = \int [V_\alpha] d\mu(\alpha)$. Suppose that $\lambda = dw - f'_\alpha(z)dz$. We want to show that $\lambda \wedge T = 0$. Let ϕ be any smooth $(1, 0)$ test form. We need to show that $\langle \lambda \wedge T, \phi \rangle = 0$. This follows since

$$\begin{aligned}
\langle \lambda \wedge T, \phi \rangle &= \int (\lambda \wedge T) \wedge \phi \\
&= \int T \wedge (\lambda \wedge \phi) \\
&= \int_{\alpha} \left(\int_{V_{\alpha}} \lambda \wedge \phi \right) d\mu(\alpha) \\
&= \int_{\alpha} 0 = 0
\end{aligned}$$

Assume next that T is weakly directed by $\mathcal{L}$. Since $\mathcal{L}$ is a lamination of $\Delta \times \mathbb{C}$ we may invoke the approximation result from the previous section. With the approximation result at hand the implication follows from Sullivan's proof of the smooth case [7]. We include the proof for the benefit of the reader.

Step 1 is to show that there exists a family of probability measures σ_{α} such that σ_{α} is supported on Γ_{α} , and a measure μ' on the α -plane, such that for all test forms ω we have that

$$T(\omega) = \int \left(\int_{\Gamma_{\alpha}} \omega d\sigma_{\alpha} \right) d\mu'.$$

Let ω be a $(1,1)$ test form and let $\lambda(z, w) = dw - f'_{\alpha}(z)dz$ for $w = f_{\alpha}(z)$. Let $\vec{v}_1(z, w) = (1, f'_{\alpha}(z))$ and let $\vec{v}_2(z, w) = (i, i \cdot f'_{\alpha}(z))$ for $w = f_{\alpha}(z)$, and define the 2-tangent field $v(z, w) = (\vec{v}_1(z, w), \vec{v}_2(z, w))$.

Switching basis we have that

$$\omega = \psi_1 dz \wedge d\bar{z} + \psi_2 dz \wedge \bar{\lambda} + \psi_3 d\bar{z} \wedge \lambda + \psi_4 \lambda \wedge \bar{\lambda}$$

for some functions ψ_i , and by assumption we have that $T(\omega) = T(\psi_1 dz \wedge d\bar{z})$. The function ψ_1 is given by $\psi_1 = \frac{1}{2i}\omega(v)$ and so we have that

$$T(\omega) = T\left(\frac{1}{2i}\omega(v)dz \wedge d\bar{z}\right).$$

On the other hand we may use T to define a linear functional L on $\mathcal{C}_0(\Delta \times \mathbb{C})$ by $L(\psi) = T(\psi dz \wedge d\bar{z})$, and so by Riesz' Representation Theorem there is a measure ν such that

$$L(\psi) = \int \psi d\nu.$$

This means that

$$T(\omega) = \int \frac{1}{2i}\omega(v)d\nu.$$

Now the measure ν disintegrates [3]: There exists a family of probability measures σ_{α} such that σ_{α} is supported on Γ_{α} , and a measure μ' on the α -plane, such that for all $\psi \in \mathcal{C}_0(\Delta \times \mathbb{C})$ we have that

$$\int \psi d\nu = \int \left(\int_{\Gamma_{\alpha}} \psi d\sigma_{\alpha} \right) d\mu'.$$

We define currents T_{α} by $T_{\alpha}(\omega) = \int_{\Gamma_{\alpha}} \frac{1}{2i}\omega(v)d\sigma_{\alpha}$, and we get that

$$T(\omega) = \int T_{\alpha}(\omega)d\mu'.$$

The next step is to show that T_α is closed for μ' -almost all α . Let $\{\omega_j\}$ be a dense set of $\mathcal{C}^1$ -smooth $(0,1)$ test forms and fix a $j \in \mathbb{N}$. Let g be a continuous function in the α -variable and extend g constantly along leaves. We want to show that

$$\int g \cdot T_\alpha(\partial\omega) d\mu' = 0$$

because this would imply that $\partial T_\alpha = 0$ for μ' -almost all α (since g is arbitrary).

By Theorem 1 there exists a sequence g_i of smooth functions such that $g_i \rightarrow g$ uniformly and in $\mathcal{C}^1$ -norm on leaves. Since T is closed we have that

$$0 = \int T_\alpha(\partial(g\omega_j)) d\mu' = \int T_\alpha(\partial g_i \wedge \omega_j) d\mu' + \int g_i \cdot T_\alpha(\partial\omega_j) d\mu'.$$

Since $T_\alpha(\partial g_i \wedge \omega) \rightarrow 0$ we get that

$$\int g \cdot T_\alpha(\partial\omega_j) d\mu' = \lim_{i \rightarrow \infty} \int g_i \cdot T_\alpha(\partial\omega_j) d\mu' = 0$$

Running through all ω_j 's we see that T_α is closed for μ' -almost all α . The only possibility then is that the measures σ_α are constant multiples of $dz \wedge d\bar{z}$, i.e. $\sigma_\alpha = \varphi(\alpha) dz \wedge d\bar{z}$ where φ is a measurable function [4]. Define $\mu := \varphi \cdot \mu'$.

5. TWO COUNTEREXAMPLES

In [2] the authors proved versions of the main theorem for real laminations in $\mathbb{R}^2$ and $\mathbb{R}^3$. In those results we added an extra slope condition on the laminations which is analogous to the estimate in Corollary 1. We give here a simple example of a lamination of curves in $\mathbb{R}^2$ where the slope condition is not satisfied. Also the conclusion of the Main Theorem fails. The analogue of Theorem 1, i.e. approximation of partially smooth functions fails as well.

For each $t \in \mathbb{R}$, we let γ_t be the curve $y = f_t(x) = (x-t)^3$ in $\mathbb{R}^2$. Clearly this gives a continuous lamination of $\mathbb{R}^2$ by curves. The curves are all tangent to the x -axis. This implies that the current of integration of the x -axis is annihilated by the one form λ which is defined by $dy - f'_t(x)dx$ on γ_t . However, this current is not an integral of currents $[\gamma_t]$. We also observe that the function $a(x, y)$ defined by $a(x, f_t(x)) = t$ cannot be approximated by $\mathcal{C}^1$ functions, because any such approximation will have to have a small derivative along the x -axis.

We can also modify this example so that we have a Riemann surface lamination in $\mathbb{C}^3$. For $t \in \mathbb{C}$, let γ_t be the complex curve $\gamma_t(s) = (z, w, \tau) = (s, (s-t)^2, (s-t)^3)$. These curves laminate $\mathbb{C}^3$ and γ_t is tangent to the z -axis at $(t, 0, 0)$. Hence the z -axis is annihilated by any continuous one forms defining the lamination. Hence the current of integration of the z -axis is weakly directed. But clearly it is not directed by the lamination. Again the function $a(z, w, \tau)$ defined by $a|_{\gamma_t} = t$ cannot be approximated by $\mathcal{C}^1$ functions.

REFERENCES

- [1] Fornæss, J. E., Sibony, N; *Harmonic currents and finite energy of laminations*, GAFA **15** (2005), 962–1003.
- [2] Fornæss, J. E., Wang, Y., Wold, E. F; *Approximation of partially smooth functions*, Proceedings of Qikeng Lu conference, June 2006, Science in China (Ser A), (2007) to appear.
- [3] Halmos, P.R.; *Measure Theory*, Van Nostrand (1950).

- [4] Lelong, P.; *Fonctions plurisubharmoniques et formes différentielle positive*, Gordon and Breach, Paris-London-New York (1968).
- [5] Mañé, P., Sad P., Sullivan D.; *On the dynamics of rational maps*, Ann.Sci.École Norm.Sup. **16** (1983), 193–217
- [6] Ślodkowski, Z; *Holomorphic motions and polynomial hulls*, Proc. Amer. Math. Soc. **111** (1991), 347–355.
- [7] Sullivan, D; *Cycles for the dynamical study of foliated manifolds and complex manifolds*, Invent. Math. **36** (1976), 225–255.
- [8] Astala, K., Martin, G. J; *Holomorphic Motions*, In Papers on Analysis, A volume dedicated to Olli Martio on the occasion of his 60th birthday. Report Univ. Jyväskylä, Jyväskylä **83** (2001), 27–40.

John Erik Fornæss
 Mathematics Department
 The University of Michigan
 East Hall, Ann Arbor, MI 48109
 USA
 fornaess@umich.edu

Yinxia Wang
 Department of Mathematics
 Henan University
 Kaifeng, 475001
 China
 yinxiaawang@gmail.com

Erlend Fornæss Wold
 Mathematisches Institut
 Universität Bern
 Sidlerstr. 5
 CH-3012 Bern
 Switzerland
 erlendfw@math.uio.no